

The C-H bond activation in 1-ethyl-3-methylimidazolium acetate-copper(ii) acetate-water-air (dioxygen) systems

Shtyrlin V., Serov N., Islamov D., Konkin A., Bukharov M., Gnezdilov O., Krivolapov D., Kataeva O., Nazmutdinova G., Wendler F.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Ionic liquid (1-ethyl-3-methylimidazolium acetate, [C2C 1im][AcO])-copper(ii) diacetate monohydrate-water-air (O₂) systems have been investigated by ¹³C NMR, EPR, spectrophotometry, HPLC, and synthetic chemistry methods at different temperatures. The C-H bond activation of [C2C1im]⁺ with the formation of the unusual dication 1,1'-diethyl-3,-'-dimethyl-2,2'-biimidazolium ([C2C1im)₂]²⁺) at 50°C and 1-ethyl-3-methyl-1H-imidazol-2(3H)-one (C2C 1imO) at 50-85°C was revealed. Two new complexes with the above compounds, [(C2C1im)₂][Cu(AcO)₄] and Cu₂(AcO)₄(C2C1imO)₂, were isolated from the systems and characterized by X-ray structural analysis. Catalytic cycles with the participation of copper(ii) acetate and dioxygen and the production of [(C2C1im)₂]²⁺ and C2C1imO have been proposed. The catalysis presumably includes the formation of the CuII(O₂)CuII active centre with μ-η²:η²-peroxide bridging in analogy with tyrosinase and catechol oxidase activity. © 2014 The Royal Society of Chemistry.

<http://dx.doi.org/10.1039/c3dt51946e>
